

# 1 Antiderivatives

## 1.1 T/F

1. (True False) Every function has an antiderivative.
2. (True False) Every continuous function has an antiderivative.
3. (True False) Every differentiable function has an antiderivative.
4. (True False) Antiderivatives are usually harder than derivatives.
5. (True False) The antiderivative of the derivative of a function is equal to the function itself.
6. (True False) The derivative of an antiderivative of a function is equal to the function itself.

## 1.2 Exercises

Find the antiderivatives of the following functions (you may need to “undo” some derivative rules):

1. 0 (the constant 0 function)
2.  $x^{-204.5}$
3.  $\frac{1+x^2+x^3}{x^7}$
4.  $\frac{1}{2\sqrt{x}}$
5.  $10^x$
6.  $\sec(x) \tan(x)$
7.  $e^{\sin(x)} \cos(x)$
8.  $x \cos(x^2)$
9.  $\sin(x) \cos(x)$
10.  $\frac{1}{4+x^2}$
11.  $\frac{\sin(x)^3 + \sin(x) \cos(x)^2}{\cos(x)^2}$
12.  $\csc(x) \cot(x)$

## 2 Riemann sums

### 2.1 T/F

1. (True False) Riemann sums always exist for any function.
2. (True False) Let  $R_n$  be the Riemann sum for a function  $f(x)$  on the interval  $[a, b]$  using  $n$  evenly spaced rectangles. The limit,  $\lim_{n \rightarrow \infty} R_n$ , if it exists, is the area under the curve of  $f(x)$  from  $a$  to  $b$ . (If the limit exists,  $f(x)$  is called *integrable* on  $[a, b]$ .)
3. (True False) Riemann sums using right endpoints generally overestimate the area of a function.
4. (True False) Riemann sums using left endpoints generally underestimate the area of a function.
5. (True False) Riemann sums are not useful because they are so difficult to compute.
6. (True False) Finite Riemann sums always exist for any function.

### 2.2 Exercises

1. Compute  $R_4$  (same meaning as above) for  $f(x) = x^2$  on  $[0, 1]$  with left endpoints. Does this overestimate or underestimate the actual area? (For the last part, a picture is helpful.)
2. Compute  $R_4$  (same meaning as above) for  $f(x) = x^2$  on  $[0, 1]$  with right endpoints. Does this overestimate or underestimate the actual area?
3. Think about when right Riemann sums will over/underestimate the area. Same with left Riemann sums.
4. Compute  $L_4$  for  $f(x) = e^{|x|}$  on  $[-1, 1]$  with right endpoints. Do it with left endpoints.
5. Let  $f(x)$  be the function defined by:

$$f(x) = \begin{cases} 0 & x \in \mathbb{Q} \\ 1 & x \notin \mathbb{Q} \end{cases}$$

Compute  $R_{1000}$  for  $f(x)$  on  $[0, 1]$  with right endpoints. Compute  $R_{1000}$  for  $f(x)$  on  $[0, \pi]$  with right endpoints.

Do you think the Riemann sum  $\lim_{n \rightarrow \infty} R_n$  exists? Why or why not?